

***What is claimed is:***

1. A method of identifying one or more markers for osteoarthritis, wherein each of said one or more markers corresponds to a gene transcript, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from one or more individuals having osteoarthritis, wherein each of said one or more transcripts is expressed by a gene that is a candidate marker for osteoarthritis; and

b) comparing the level of each of said one or more gene transcripts from said step a) with the level of each of said one or more genes transcripts in blood obtained from one or more individuals not having osteoarthritis,

wherein those compared transcripts which display differing levels in the comparison of step b) are identified as being markers for osteoarthritis.

2. A method of identifying one or more markers for osteoarthritis, wherein each of said one or more markers corresponds to a gene transcript, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from one or more individuals having osteoarthritis, wherein each of said one or more transcripts is expressed by a gene that is a candidate marker for osteoarthritis; and

b) comparing the level of each of said one or more gene transcripts from said step a) with the level of each of said one or more genes transcripts in blood obtained from one or more individuals having osteoarthritis,

wherein those compared transcripts which display the same levels in the comparison of step b) are identified as being markers for osteoarthritis.

3. A method of identifying one or more markers of a stage of osteoarthritis progression or regression, wherein each of said one or more markers corresponds to a gene transcript, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from one or more individuals having a stage of osteoarthritis, wherein said one or more individuals are at the same progressive or regressive stage of osteoarthritis, and wherein each of said one or more transcripts is expressed by a gene that is a candidate marker for determining the stage of progression or regression of osteoarthritis, and;

b) comparing the level of each of said one or more gene transcripts from said step a) with

the level of each of said one or more genes transcripts in blood obtained from one or more individuals who are at a progressive or regressive stage of osteoarthritis distinct from that of said one or more individuals of step a),

wherein those compared transcripts which display differing levels in the comparison of step b) are identified as being markers for the stage of progression or regression of osteoarthritis.

4. A method of identifying one or more markers of a stage of osteoarthritis progression or regression, wherein each of said one or more markers corresponds to a gene transcript, comprising the steps of:

- a) determining the level of one or more gene transcripts expressed in blood obtained from one or more individuals having a stage of osteoarthritis, wherein said one or more individuals are at the same progressive or regressive stage of osteoarthritis, and wherein each of said one or more transcripts is expressed by a gene that is a candidate marker for determining the stage of progression or regression of osteoarthritis, and;

- b) comparing the level of each of said one or more gene transcripts from said step a) with the level of each of said one or more genes transcripts in blood obtained from one or more individuals who are at a progressive or regressive stage of osteoarthritis identical to that of said one or more individuals of step a),

wherein those compared transcripts which display the same levels in the comparison of step b) are identified as being markers for the stage of progression or regression of osteoarthritis.

5. The method of any one of claims 1-4, wherein each of said one or more markers identifies one or more transcripts of one or more non immune response genes.
6. The method of any one of claims 1-4, wherein each of said one or more markers identifies a transcript of a gene expressed by non-blood tissue.
7. The method of any one of claims 1-4, wherein each of said one or more markers identifies a transcript of a gene expressed by non-lymphoid tissue.

8. The method of any one of claims 1-4, wherein each of said one or more markers identifies a transcript of a gene selected from the group consisting of the genes listed in Table 3O and Table 3AB.

9. A method of diagnosing, or prognosing osteoarthritis in an individual, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from said individual, wherein said one or more gene transcripts corresponds to said one or more markers of claim 1 and claim 2, and

b) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from one or more individuals not having osteoarthritis,

wherein detecting a difference in the levels of each of said one or more gene transcripts in the comparison of step b) is indicative of osteoarthritis in the individual of step a).

10. A method of diagnosing or prognosing osteoarthritis in an individual, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from said individual, wherein said one or more gene transcripts correspond to said one or more markers of claim 1 and claim 2, and

b) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from one or more individuals having osteoarthritis,

wherein detecting the same levels of each of said one or more gene transcripts in the comparison of step b) is indicative of osteoarthritis in the individual of step a).

11. A method of determining a stage of disease progression or regression in an individual having osteoarthritis, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from said individual having osteoarthritis, wherein said one or more gene transcripts correspond to said one or more markers of claim 3 and claim 4, and

b) comparing the level of each of said one or more gene transcripts in said blood

according to step a) with the level of each of said one or more gene transcripts in blood obtained from one or more individuals who each have been diagnosed as being at the same progressive or regressive stage of osteoarthritis,

wherein the comparison from step b) allows the determination of the stage of osteoarthritis progression or regression in the individual of step a).

12. A method of diagnosing or prognosing osteoarthritis in an individual, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from said individual, wherein said one or more gene transcripts correspond to said one or more markers of claim 1 and claim 2, and

b) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from one or more individuals having osteoarthritis,

c) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from one or more individuals not having osteoarthritis,

d) determining whether the level of said one or more gene transcripts of step a) classify with the levels of said transcripts in step b) as compared with the levels of said transcripts in step c)

wherein said determination is indicative of said individual of step a) having osteoarthritis.

13. A method of determining a stage of disease progression or regression in an individual having osteoarthritis, comprising the steps of:

a) determining the level of one or more gene transcripts expressed in blood obtained from said individual having said stage of osteoarthritis, wherein said one or more gene transcripts correspond to the markers of claim 3 and claim 4, and

b) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from one or more individuals having said stage of osteoarthritis,

c) comparing the level of each of said one or more gene transcripts in said blood according to step a) with the level of each of said one or more gene transcripts in blood from

one or more individuals not having said stage of osteoarthritis,

d) determining whether the level of said one or more gene transcripts of step a) classify with the levels of said transcripts in step b) as compared with levels of said transcripts in step c)

wherein said determination is indicative of said individual of step a) having said stage of osteoarthritis.

14. The method of any one of claims 9-13, wherein said one or more gene transcripts are transcribed from one or more genes selected from the group consisting of Table 3O and Table 3AB.

15. The method of any one of claims 1-4 and 9-13, wherein said one or more gene transcripts are transcribed from one or more genes selected from the group consisting of : a) non-immune response genes, b) genes expressed by non blood tissue, and c) genes expressed by non lymphoid tissue.

16. The method of any one of claims 1-4 and 9-13, wherein said blood comprises a blood sample obtained from said one or more individuals.

17. The method of claim 16, wherein said blood sample consists of whole blood.

18. The method of claim 16, wherein said blood sample consists of a drop of blood.

19. The method of claim 16, wherein said blood sample consists of blood that has been lysed.

20. The method of claim 16, further comprising the step of isolating RNA from said blood samples.

21. The method of any one of claims 1-4 and 9-13, wherein the step of determining the level of each of said one or more gene transcripts comprises quantitative RT-PCR (QRT-PCR), wherein said one or more transcripts are from step a) and/or step b) of claims 1-4 and 9-13.

22. The method of claim 21, wherein said QRT-PCR comprises primers which hybridize to said one or more transcripts or the complement thereof, wherein said one or more transcripts are from step a) and/or step b) of claims 1-4 and 9-13.

23. The method of claim 22, wherein said primers are 15-25 nucleotides in length.
24. The method of claim 22, wherein said primers hybridize to one or more genes selected from the group consisting of the genes listed in Table 3O and Table 3AB, or the complement thereof.
- 5 25. The method of any one of claims 1-4 and 9-13, wherein the step of determining the level of each of said one or more gene transcripts comprises hybridizing a first plurality of isolated nucleic acid molecules that correspond to said one or more transcripts, to an array comprising a second plurality of isolated nucleic acid molecules.
26. The method of claim 25, wherein said first plurality of isolated nucleic acid molecules  
10 comprises RNA, DNA, cDNA, PCR products or ESTs.
27. The method of claim 25, wherein said array comprises a plurality of isolated nucleic acid molecules comprising RNA, DNA, cDNA, PCR products or ESTs.
28. The method of claim 27, wherein said array comprises two or more of the markers of claim 1.
- 15 29. The method of claim 27, wherein said array comprises two or more of the markers of claim 2.
30. The method of claim 27, wherein said array comprises two or more of the markers of claim 3.
31. The method of claim 27, wherein said array comprises two or more of the markers of claim  
20 4.
32. The method of claim 27, wherein said array comprises a plurality of nucleic acid molecules that correspond to genes of the human genome.
33. The method of claim 27, wherein said array comprises a plurality of nucleic acid molecules that correspond to two or more genes selected from the group consisting of the genes listed in  
25 Table 3O and Table 3AB.

34. A plurality of isolated nucleic acid molecules that correspond to two or more of the markers of claim 1.
35. A plurality of isolated nucleic acid molecules that correspond to two or more of the markers of claim 2.
- 5 36. A plurality of isolated nucleic acid molecules that correspond to two or more of the markers of claim 3.
37. A plurality of isolated nucleic acid molecules that correspond to two or more of the markers of claim 4.
38. The method of claim 26, wherein said ESTs comprise at least 100 nucleotides in length.
- 10 39. An array consisting essentially of the plurality of nucleic acid molecules of claim 34.
40. An array consisting essentially of the plurality of nucleic acid molecules of claim 35.
41. An array consisting essentially of the plurality of nucleic acid molecules of claim 36.
42. An array consisting essentially of the plurality of nucleic acid molecules of claim 37.
- 15 43. A kit for diagnosing or prognosing osteoarthritis comprising: a) two gene-specific priming means designed to produce double stranded DNA complementary to a gene that corresponds to a marker selected from the group consisting of the markers of Claim 1, Claim 2, Claim 3 and Claim 4; wherein said first priming means contains a sequence which can hybridize to RNA, cDNA or an EST complementary to said gene to create an extension product and said second priming means capable of hybridizing to said extension product; b) an enzyme with reverse transcriptase activity c) an enzyme with thermostable DNA polymerase activity and  
20 d) a labeling means; wherein said primers are used to detect the quantitative expression levels of said gene in a test subject.
- 25 44. A kit for monitoring a course of therapeutic treatment of osteoarthritis, comprising a) two gene-specific priming means designed to produce double stranded DNA complementary to a gene that corresponds to a marker selected from the group consisting of the markers of Claim

1, Claim 2, Claim 3 and Claim 4; wherein said first priming means contains a sequence which can hybridize to RNA, cDNA or an EST complementary to said gene to create an extension product and said second priming means capable of hybridizing to said extension product; b) an enzyme with reverse transcriptase activity c) an enzyme with thermostable DNA polymerase activity and d) a labeling means; wherein said primers are used to detect the quantitative expression levels of said gene in a test subject.

45. A kit for monitoring progression or regression of osteoarthritis, comprising: a) two gene-specific priming means designed to produce double stranded DNA complementary to a gene that corresponds to a marker selected from the group consisting of the markers of Claim 1, Claim 2, Claim 3 and Claim 4; wherein said first priming means contains a sequence which can hybridize to RNA, cDNA or an EST complementary to said gene to create an extension product and said second priming means capable of hybridizing to said extension product; b) an enzyme with reverse transcriptase activity c) an enzyme with thermostable DNA polymerase activity and d) a labeling means; wherein said primers are used to detect the quantitative expression levels of said gene in a test subject.

46. The kit of any one of claims 43-45 wherein said gene-specific priming means are complementary to a gene selected from the group consisting of the genes listed in Table 3O and Table 3AB.

47. A plurality of nucleic acid molecules that identify or correspond to two or more genes selected from the group consisting of the genes listed in Table 3O and Table 3AB.

48. The method of claim 27, wherein said ESTs comprise at least 100 nucleotides in length.